

OpenType math font Fira

Herbert Voß

March 21, 2019

Contents

| | |
|-------------------------------------|----------|
| 1 Usage | 1 |
| 2 The default regular weight | 2 |
| 2.1 Version normal | 2 |
| 2.2 Version bold | 2 |
| 3 Examples | 2 |
| 3.1 Digits | 2 |
| 3.2 Alphabets | 3 |
| 3.3 Equations test | 4 |

Abstract

The math font FIRA is derived from the Fira Sans and Fira Go sans serif. There are several math versions available (<https://github.com/Stone-Zeng/FiraMath/>) but only the regular version has from todays update all symbols.

1 Usage

```
\usepackage[<options>]{firamath-otf}
```

Optional arguments are

fakebold Use faked bold symbols

usefilenames Use filenames for the fonts instead of the symbolic font names

The package itself loads by default

```
\RequirePackage{ifxetex,ifluatex,xkeyval,textcomp}  
\RequirePackage{unicode-math}
```

2 The default regular weight

2.1 Version normal

$$\begin{aligned}\frac{\partial \rho}{\partial t} + \text{div}(\rho \vec{v}) &= 0 \\ \rho \frac{\partial \vec{v}}{\partial t} + (\rho \vec{v} \cdot \nabla) \vec{v} &= \vec{f}_0 + \text{div} \mathbf{T} = \vec{f}_0 - \text{grad } p + \text{div} \mathbf{T}' \\ \rho T \frac{ds}{dt} &= \rho \frac{de}{dt} - \frac{p}{\rho} \frac{d\rho}{dt} = -\text{div} \vec{q} + \mathbf{T}' : \mathbf{D}\end{aligned}\quad (1)$$

$$\begin{aligned}\frac{\partial}{\partial t} \iiint \rho d^3V + \oint \rho (\vec{v} \cdot \vec{v} \vec{n}) d^2A &= 0 \\ \frac{\partial}{\partial t} \iiint \rho \vec{v} d^3V + \oint \rho \vec{v} (\vec{v} \cdot \vec{n}) d^2A &= \iiint f_0 d^3V + \oint \vec{n} \cdot \mathbf{T} d^2A\end{aligned}\quad (2)$$

$$\begin{aligned}\frac{\partial}{\partial t} \iiint \left(\frac{1}{2} v^2 + e \right) \rho d^3V + \oint \left(\frac{1}{2} v^2 + e \right) \rho (\vec{v} \cdot \vec{n}) d^2A &= \\ - \oint (\vec{q} \cdot \vec{v} \vec{n}) d^2A + \iiint (\vec{v} \cdot \vec{f}_0) d^3V + \oint (\vec{v} \cdot \vec{n} \mathbf{T}) d^2A.\end{aligned}\quad (3)$$

2.2 Version bold

The bold characters are created with the optional argument `fakebold` which loads the package `xfakebold` which writes some information into the created PDF to get bold characters. For more informations see the documentation of `xfakebold`.

$$\begin{aligned}\frac{\partial}{\partial t} \iiint \rho d^3V + \oint \rho (\vec{v} \cdot \vec{v} \vec{n}) d^2A &= 0 \\ \frac{\partial}{\partial t} \iiint \rho \vec{v} d^3V + \oint \rho \vec{v} (\vec{v} \cdot \vec{n}) d^2A &= \iiint f_0 d^3V + \oint \vec{n} \cdot \mathbf{T} d^2A\end{aligned}\quad (4)$$

$$\begin{aligned}\frac{\partial}{\partial t} \iiint \left(\frac{1}{2} v^2 + e \right) \rho d^3V + \oint \left(\frac{1}{2} v^2 + e \right) \rho (\vec{v} \cdot \vec{n}) d^2A &= \\ - \oint (\vec{q} \cdot \vec{v} \vec{n}) d^2A + \iiint (\vec{v} \cdot \vec{f}_0) d^3V + \oint (\vec{v} \cdot \vec{n} \mathbf{T}) d^2A.\end{aligned}\quad (5)$$

3 Examples

3.1 Digits

- Digits:

0123456789

- Proportional digits: 0123456789
- Bold digits (`\symbf`): **0123456789**
- Bold proportional digits (`\symbf`): **0123456789**

3.2 Alphabets

- Latin letters (`\mathnormal`):
ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz
- Latin upright letters (`\symup`):
ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz
- Latin typewriter letters (`\symtt`):
ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz
- Latin bold letters (`\symbf`):
ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz
- Latin bold upright letters (`\symbfup`):
ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz
- Latin blackboard letters (`\symbb`):
ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz
- Greek letters:
ΑΒΓΔΕΖΗΘΙΚΛΜΝΞΟΠΡΣΤΥΦΧΨΩαβγδεζηθικιλμνξοπρρςςτυφρχψω
- Greek upright letters (`\symup`):
ΑΒΓΔΕΖΗΘΙΚΛΜΝΞΟΠΡΣΤΥΦΧΨΩαβγδεζηθικιλμνξοπρρςςτυφρχψω
- Greek bold letters (`\symbf`):
ΑΒΓΔΕΖΗΘΙΚΛΜΝΞΟΠΡΣΤΥΦΧΨΩαβγδεζηθικιλμνξοπρρςςτυφρχψω
- Greek bold upright letters (`\symbfup`):
ΑΒΓΔΕΖΗΘΙΚΛΜΝΞΟΠΡΣΤΥΦΧΨΩαβγδεζηθικιλμνξοπρρςςτυφρχψω
- Dotless letters:
ı + j + ı + j
- Hebrew *א + ב + ג + ד*
- Ligature (text):
ff fi fl ffi ffl
- Non-ligature (math):
ff fi fl ffi ffl + ff fi fl ffi ffl + ff fi fl ffi ffl
- Miscellaneous:
ħ + ħ + Å
∀x > x₀, ∃δ, δ ∈ ∅

3.3 Equations test

- Basic:
 $1 + 2 - 3 \times 4 \div 5 \pm 6 \mp 7 + 8 = -a \oplus b \otimes c$
- Binary relations $x + - \oplus \otimes \odot \oslash \cdot \cdot \times \div y$
- Set theory $A \cap B \cup C \cap D \sqcup R \uplus k \cup l \uplus m$
 $A \subset B \supset C \subseteq D \supseteq E \quad F \setminus G + A \subset B \supset C \subseteq D \supseteq E$
 $\mathbb{C}_U A \cup \mathbb{C}_C C \subset \mathbb{C}_U A \cup \mathbb{C}_C C \in R \in Q \ni Z \ni N$
- Superscript and subscript:
 $2^2 + 2^{2^2} + 2^{2^{2^2}} + 2^{2^2} + x_a + x_{a_i} + x_{a_{i_1}}$
- Arrows:
 $x \leftarrow y \rightarrow z \leftrightarrow w \nleftrightarrow y \rightharpoonup z \leftrightsquigarrow w \Leftarrow a \Rightarrow b \Leftrightarrow c \quad a = b \quad c$
 $x \uparrow y \downarrow z \Downarrow w \Uparrow a \Downarrow b \Updownarrow c$
 $p \curvearrowright p \nearrow p \searrow p \swarrow p \nwarrow p \nearrow p \searrow p \nrightarrow p \not\rightarrow p$
 $x \leftarrow x \leftarrow x \uparrow x \uparrow x \rightarrow x \rightarrow x \downarrow x \downarrow x$
 $A \longleftarrow B \longrightarrow C \longleftrightarrow D \Longleftarrow E \Longrightarrow F \Longleftrightarrow G$
 $X \leftrightsquigarrow Y \mapsto Z \Uparrow W \Downarrow P \Leftarrow S \Rightarrow R$
 $M \longleftrightarrow N \mapsto O \Longleftarrow K \Longrightarrow L$
 $f \rightrightarrows f \Updownarrow f \leftrightsquigarrow f \Uparrow g \rightrightarrows g \Uparrow g \Leftarrow g \Downarrow h \rightrightarrows h \Longleftarrow p \rightrightarrows p \Leftarrow p \Uparrow p \Uparrow p$
- Math accents:

x̂ x̃ x̄ x̅ x̆ ẋ ẍ x̉ x̊ x̋ x̌ x̍ x̎ x̏ x̐ x̑ x̒ x̓ x̔ x̕ x̖ x̗ x̘ x̙ x̚ x̛ x̜ x̝ x̞ x̟ x̠ x̡ x̢ x̣ x̤ x̥ x̦ x̧ x̨ x̩ x̪ x̫ x̬ x̭ x̮ x̯ x̰ x̱ x̲ x̳ x̴ x̵ x̶ x̷ x̸ x̹ x̺ x̻ x̼ x̽ x̾ x̿ x̺ x̻ x̼ x̽ x̾ x̿ x̺ x̻ x̼ x̽ x̾ x̿

- Integral:

$$\int_0^{\pi} \sin x \, dx = \int_0^{\pi} \sin x \, dx = \cos 0 - \cos \pi = 2$$

$$\int_{-\infty}^{+\infty} dz \iint_{-\infty}^{+\infty} d^2y \iiint_{-\infty}^{+\infty} d^3x \iiiii_{-\infty}^{+\infty} d^4p$$

$$\oint dr \oint d\theta \oint d\varphi$$

$$\int_0^{\pi} \sin x \, dx = \int_0^{\pi} \sin x \, dx = \cos 0 - \cos \pi + C$$

$$\int_{-\infty}^{+\infty} dz \iint_{-\infty}^{+\infty} d^2 y \iiint_{-\infty}^{+\infty} d^3 x \iiint_{-\infty}^{+\infty} d^4 p$$

$$\oint dr \oint d\theta \oint d\varphi$$

- Huge operators:

$$\int_0^\infty \int_0^\infty \sum_{i=1}^\infty \prod_{j=i}^\infty \prod_{k=i}^\infty$$

$$\sum_{i=1}^\infty \frac{1}{x^i} = \frac{1}{1-x} \quad \prod_{i=1}^\infty \frac{1}{x^i} = x^{-n(n+1)/2} \quad \prod_{i=1}^\infty \frac{1}{x^i} = ?$$

- Huge operators (inline):

$$\int_0^\infty \int_0^\infty \iint dx \iiint dy \iiiii dp \oint dr \oslash d\theta \oslash\oslash d\varphi \sum_{i=1}^\infty \prod_{j=i}^\infty \prod_{i=i}^\infty$$

- Huge operators (inline):

$$\int_0^\infty \int_0^\infty \iint dx \iiint dy \iiiii dp \oint dr \oslash d\theta \oslash d\varphi \sum_{i=1}^\infty \prod_{j=i}^\infty \prod_{i=i}^\infty$$

- Fraction:

$$\frac{1}{2} + \frac{1}{\frac{2}{3} + 4} + \frac{\frac{1}{2} + 3}{4}$$

- Fraction (inline):

$$\frac{1}{2} + \frac{1g}{2} + \frac{1}{\frac{2}{3} + 4} + \frac{\frac{1}{2} + 3}{4}$$

- Radical:

$$\sqrt{2} + \sqrt{2^2} + \sqrt{1 + \sqrt{2}} + \sqrt{1 + \sqrt{1 + \sqrt{3}}} + \sqrt{\sqrt{\sqrt{\sqrt{2}}}} + \sqrt{\frac{1}{2}}$$

$$\sqrt[3]{2} + \sqrt[3]{2^2} + \sqrt[3]{1 + \sqrt[3]{2}} + \sqrt[3]{1 + \sqrt[3]{1 + \sqrt[3]{3}}} + \sqrt[3]{\sqrt[3]{3}\sqrt[3]{\sqrt[3]{2}}} + \sqrt[3]{\frac{1}{2}}$$

$$\sqrt[4]{2} + \sqrt[4]{2^2} + \sqrt[4]{1 + \sqrt[4]{2}} + \sqrt[4]{1 + \sqrt[4]{1 + \sqrt[4]{3}}} + \sqrt[4]{\sqrt[4]{4}\sqrt[4]{\sqrt[4]{2}}} + \sqrt[4]{\frac{1}{2}}$$

$$\sqrt[x]{y} + \sqrt{\sqrt[x]{y}} + \sqrt{\sqrt{\sqrt[x]{y}}} + \sqrt{\frac{1}{2}} + \sqrt{\frac{x}{z}y} + \sqrt[3]{\frac{x}{z}y} + \sqrt[4]{\frac{x}{z}y} + \sqrt[x]{\frac{x}{z}y} + \sqrt{\frac{x}{w}y} + \sqrt[3]{\frac{x}{w}y} + \sqrt[4]{\frac{x}{w}y} + \sqrt[x]{\frac{x}{w}y}$$

- Bra-kets:

$$\langle x| + |x\rangle + \langle \alpha|\beta\rangle + |\alpha^2\rangle\langle\beta^2| + \left\langle \frac{1}{2} \right| + \left| \frac{1}{2} \right\rangle + \left\langle \frac{1}{2} \middle| \frac{1}{2} \right\rangle + \left| \frac{1}{2} \right\rangle \left\langle \frac{1}{2} \right| + \left\langle \frac{a^2}{b^2} \right| + \left| \frac{e^{x^2}}{e^{y^2}} \right\rangle$$

$$\langle | \rangle \quad \langle | \rangle \quad \langle | \rangle \quad \langle | \rangle \quad \langle | \rangle \quad \langle | \rangle \quad \langle | \rangle \quad \langle | \rangle \quad \langle | \rangle \quad \langle | \rangle \quad \langle | \rangle$$

- Matrices:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} + \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

$$\begin{pmatrix} a & b & c & d \\ x & y & z & w \end{pmatrix} \quad \begin{bmatrix} a & b & c & d \\ x & y & z & w \end{bmatrix} \quad \left\{ \begin{matrix} a & b & c & d \\ x & y & z & w \end{matrix} \right\} \quad \left| \begin{matrix} a & b & c & d \\ x & y & z & w \end{matrix} \right| \quad \left\| \begin{matrix} a & b & c & d \\ x & y & z & w \end{matrix} \right\|$$

$$\begin{pmatrix} a & b & c & d \\ k & l & m & n \\ x & y & z & w \end{pmatrix} \quad \begin{bmatrix} a & b & c & d \\ k & l & m & n \\ x & y & z & w \end{bmatrix} \quad \left\{ \begin{matrix} a & b & c & d \\ k & l & m & n \\ x & y & z & w \end{matrix} \right\} \quad \left| \begin{matrix} a & b & c & d \\ k & l & m & n \\ x & y & z & w \end{matrix} \right| \quad \left\| \begin{matrix} a & b & c & d \\ k & l & m & n \\ x & y & z & w \end{matrix} \right\|$$

$$\begin{pmatrix} a & b & c & d \\ k & l & m & n \\ p & q & s & t \\ x & y & z & w \end{pmatrix} \quad \begin{bmatrix} a & b & c & d \\ k & l & m & n \\ p & q & s & t \\ x & y & z & w \end{bmatrix} \quad \left\{ \begin{matrix} a & b & c & d \\ k & l & m & n \\ p & q & s & t \\ x & y & z & w \end{matrix} \right\} \quad \left| \begin{matrix} a & b & c & d \\ k & l & m & n \\ p & q & s & t \\ x & y & z & w \end{matrix} \right| \quad \left\| \begin{matrix} a & b & c & d \\ k & l & m & n \\ p & q & s & t \\ x & y & z & w \end{matrix} \right\|$$

- Nabras:

$$\nabla x + \nabla f + \nabla \cdot \mathbf{u} + \nabla \times \mathbf{v}$$

$$\nabla \quad \nabla \quad \nabla \quad \nabla; \quad \tilde{\nabla} \quad \tilde{\nabla} \quad \tilde{\nabla} \quad \tilde{\nabla}$$

- Over-/underline and over-/underbraces

$$\overline{b} \quad \overline{ab} \quad \overline{abc} \quad \overline{abcd} \quad \overline{abcde} \quad \overline{a+b+c} \quad \overline{x_1, x_2, \dots, x_n}$$

$$\frown \quad \frown \overline{b} \quad \frown \overline{ab} \quad \frown \overline{abc} \quad \frown \overline{abcd} \quad \frown \overline{abcde} \quad \frown \overline{a+b+c} \quad \frown \overbrace{x_1, x_2, \dots, x_n}^n$$

$$\neg \quad \neg \overline{b} \quad \neg \overline{ab} \quad \neg \overline{abc} \quad \neg \overline{abcd} \quad \neg \overline{abcde} \quad \neg \overline{a+b+c} \quad \neg \overbrace{x_1, x_2, \dots, x_n}^n$$

$$\smile \quad \smile \overline{b} \quad \smile \overline{ab} \quad \smile \overline{abc} \quad \smile \overline{abcd} \quad \smile \overline{abcde} \quad \smile \overline{a+b+c} \quad \smile \overbrace{x_1, x_2, \dots, x_n}^n$$

$$\underline{\underline{b}} \quad \underline{\underline{ab}} \quad \underline{\underline{abc}} \quad \underline{\underline{abcd}} \quad \underline{\underline{abcde}} \quad \underline{\underline{a+b+c}} \quad \underline{\underline{x_1, x_2, \dots, x_n}}$$

$$\underbrace{\underline{b}} \quad \underbrace{\underline{ab}} \quad \underbrace{\underline{abc}} \quad \underbrace{\underline{abcd}} \quad \underbrace{\underline{abcde}} \quad \underbrace{\underline{a+b+c}} \quad \underbrace{\underline{x_1, x_2, \dots, x_n}}_n$$

$$\underbrace{\underline{\underline{b}}} \quad \underbrace{\underline{\underline{ab}}} \quad \underbrace{\underline{\underline{abc}}} \quad \underbrace{\underline{\underline{abcd}}} \quad \underbrace{\underline{\underline{abcde}}} \quad \underbrace{\underline{\underline{a+b+c}}} \quad \underbrace{\underline{\underline{x_1, x_2, \dots, x_n}}}_n$$

- Primes

$$x'x''x'''x''''x^{(5)}x^{(6)}x^{(7)}x^{(8)}x^{(9)}x^{(10)}$$

$$x'x''x'''x^{(4)}$$

$$x'x''x'''x^{(4)}$$

$$\lim_{x \rightarrow \infty} \frac{1}{x^2} = 0$$

$$\frac{\partial y(x)}{\partial x} = \frac{dy(x)}{dx} = y'(x)$$

$$\frac{\partial y(x)}{\partial x} = \frac{dy(x)}{dx} = y'(x)$$